

Oblique triangular antiferromagnetic phase in $\text{CsCu}_{1-x}\text{Co}_x\text{Cl}_3$

T. Ono, T. Kato*, H. Tanaka, A. Hoser[†], N. Stüßer[†] and U. Schotte[†]

Department of Physics, Tokyo Institute of Technology, 152-8551 Tokyo, Japan

[†]*Hahn-Meitner-Institut, Glienicker Straße 100, D-14109 Berlin, Germany*

(Received February 1, 2008)

The spin- $\frac{1}{2}$ stacked triangular antiferromagnet $\text{CsCu}_{1-x}\text{Co}_x\text{Cl}_3$ with $0.015 < x < 0.032$ undergoes two phase transitions at zero field. The low-temperature phase is produced by the small amount of Co^{2+} doping. In order to investigate the magnetic structures of the two ordered phases, the neutron elastic scattering experiments have been carried out for the sample with $x \approx 0.03$. It is found that the intermediate phase is identical to the ordered phase of CsCuCl_3 , and that the low-temperature phase is an oblique triangular antiferromagnetic phase in which the spins form a triangular structure in a plane tilted from the basal plane. The tilting angle which is 42° at $T = 1.6$ K decreases with increasing temperature, and becomes zero at $T_{\text{N}2} = 7.2$ K. An off-diagonal exchange term is proposed as the origin of the oblique phase.

§1. Introduction

Among the group of antiferromagnets which have a hexagonal ABX_3 triangular structure, many experimental and theoretical investigations for CsCuCl_3 have been done by several authors,¹⁻⁴⁾ due to the remarkable features of this compound. Since the Cu^{2+} ion is Jahn-Teller active, CsCuCl_3 undergoes a structural phase transition at $T_t = 423$ K, where Jahn-Teller distorted octahedra CuCl_6 order so that their longest axes form a helix along the c -axis with a repeatlength of six.⁵⁻⁷⁾ The low-symmetric crystal structure produces the antisymmetric interaction of the Dzyaloshinsky-Moriya (D-M) type between the adjacent spins in the chain with the D -vector parallel to the c -direction. The exchange interactions along and between the chains are ferromagnetic and antiferromagnetic, respectively, and their values have been evaluated as $J_0/k_B = 28$ K and $J_1/k_B = -4.9$ K^{8,9)} in the definition of $\mathcal{H} = -\sum_{\langle i,j \rangle} 2J_{ij} S_i \cdot S_j$. The magnetic phase transition occurs at $T_N = 10.5$ K.^{10,11)} In the ordered state, spins lie in the basal plane and form the 120° -structure, while along the c -direction, a long period (about 71 triangular layers) helical incommensurate arrangement is realized, due to the competition between the ferromagnetic interaction and the D-M interaction along the chain.

Since the magnitude of spin on Cu^{2+} is $S = \frac{1}{2}$, quantum fluctuations are important when one describes the magnetic behavior in the magnetic field.¹²⁾ Particularly, the magnetization process with a strong field often reveals quantum effects as macroscopic phenomena. Using quantum Monte Carlo calculations or spin wave theory, for the 2D triangular antiferromagnet (TAF) in the isotropic limit, it is predicted that the magnetization plateau appears at around one third of the saturation value M_s .^{13,14)} According to these predictions, when

the external field $\mathbf{H}$ is applied parallel to the c -axis, 2D TAF undergoes a successive phase transition as shown in Fig. 1 (b) low-field coplanar, (c) collinear and (d) high-field coplanar structure in that order. In the field region where the magnetization plateau appears, the spin structure corresponds to (c) structure. For CsCuCl_3 , in-




Fig. 1. Spin structures of 2D TAF in the magnetic field. (a) Umbrella-type structure, (b) low-field coplanar structure, (c) collinear structure and (d) high-field coplanar structure.

stead of a plateau, there is a small magnetization jump around $M_s/3$, when the field is applied parallel to the c -axis. This fact is understood as the competition between weak planar anisotropy and quantum fluctuation, i.e., Fig. 1 (a) structure is stabilized by the planar anisotropy up to $H_c = 12.5$ T $> H_s/3$ and Fig. 1 (d) structure is stabilized by quantum fluctuation above $H > H_c$ with skipping (b) and (c) structures. The existence of the plateau at $M = M_s/3$ is strongly depend on the magnetic anisotropies. Recently, Honecker *et al.*¹⁵⁾ argued that the proper combination for the interchain interaction $|J_1|/J_0$ and XXZ anisotropy $\Delta = J^z/J^{xy}$ allows the existence of the plateau at $M = M_s/3$, even in the region of $\Delta < 1$ (small planar anisotropy). Here J^α ($\alpha = z$ and xy) are anisotropic exchange constants defined by $\mathcal{H} = -2 \sum_i [J_{i,i+1}^{zz} S_i^z S_{i+1}^z + \frac{1}{2} J_{i,i+1}^{xy} (S_i^+ S_{i+1}^- + S_i^- S_{i+1}^+)]$. Therefore, it is very interesting to study how the phase transition changes, when the macroscopic anisotropy of CsCuCl_3 is controlled. Because of this motivation, we

* Present address: Faculty of Education, Chiba University, 263-8522 Chiba, Japan.

have synthesized the mixture system $\text{CsCu}_{1-x}\text{Co}_x\text{Cl}_3$ with few percents of Co^{2+} ion in the concentration x , since CsCoCl_3 is known as a pseudospin- $\frac{1}{2}$ TAF with easy axis anisotropy.¹⁷⁾ In our previous paper,¹⁸⁾ we reported the results of the magnetic measurements for $\text{CsCu}_{1-x}\text{Co}_x\text{Cl}_3$ with $x < 0.032$ and the magnetic phase diagram for temperature versus magnetic field up to 7 T. Figure 2 shows the magnetic phase diagram for the external field versus temperature for the sample with $x = 0.032$ in the field parallel to the c -axis. Before




Fig. 2. The magnetic phase diagram of $\text{CsCu}_{1-x}\text{Co}_x\text{Cl}_3$ with $x = 0.32$ for the magnetic field parallel to the c -axis.

the magnetization measurements, our first speculation was that when Co^{2+} ions are doped to reduce the planar anisotropy Fig.1 (a) structure is realized in the ground state at zero field, and (b) structure which benefits quantum fluctuation might appear in the field region $H \leq H_s/3$.

On the other hand, the experimental results of the magnetization measurements indicate that the spin structure of phase II is stabilized with the help of the axial anisotropy. With decreasing Co^{2+} concentration x , the area of phase II is reduced and then disappears when $x \leq 0.005$. Thus it is reasonable to infer that, the spin structure in the intermediate phase I is identical to that of CsCuCl_3 . However, within the magnetic measurements the spin structure of the intermediate phase and low-temperature phase cannot be determined. Therefore, in order to determine the spin structure of these two phases, we have performed neutron scattering experiments for the sample with $x \approx 0.03$.

§2. Equipment

Single crystals of $\text{CsCu}_{1-x}\text{Co}_x\text{Cl}_3$ were prepared by the Bridgman method from the melt of the single crystals of CsCuCl_3 and CsCoCl_3 . The details of the sample preparation are described in the previous paper.¹⁸⁾

Neutron scattering experiments were performed at the BER II Research Reactor of the Hahn-Meitner-Institute using the triple axis spectrometer E1 operated in the double-axis mode. The incident neutron energy was fixed at $E_i = 13.9 \text{ meV}$ by Bragg reflection at the py-

rolytic graphite monochromator, and the horizontal collimations are chosen as $40'-40'-40'$ -open. A single crystal of $\text{CsCu}_{1-x}\text{Co}_x\text{Cl}_3$ with $x \approx 0.03$ was used. We label this sample $\text{CsCu}_{0.97}\text{Co}_{0.03}\text{Cl}_3$. The sample has the shape of triangular prism 5 mm in width and 15 mm in length. The sample was mounted in the cryostat with its $[1, 1, 0]$ and $[0, 0, 1]$ axes in the scattering plane.

§3. Results

From the previous magnetic measurements, for the sample with $\text{CsCu}_{0.97}\text{Co}_{0.03}\text{Cl}_3$, undergoes phase transitions at $T_{N1} \approx 9.5 \text{ K}$ and $T_{N2} \approx 7.2 \text{ K}$. First, we confirmed that the crystal structure of the present system is identical to that of pure CsCuCl_3 , i.e., the nuclear Bragg peaks were observed at (00ℓ) with $\ell = 0 \sim 6$, which are indicative of the six-fold helix. From the nuclear Bragg peaks at $(1, 1, 0)$ and $(0, 0, 6)$, the lattice constants at helium temperature were determined as $a = 7.137 \text{ \AA}$ and $c = 18.036 \text{ \AA}$.

Figure 3 shows the scans along $Q = (1/3, 1/3, \zeta)$ and $(1/3, 1/3, 6 + \zeta)$ at $T = 1.6 \text{ K}$ in the phase II. The magnetic Bragg scattering is composed of one central peak at $\zeta = 0$ and two side peaks at $\zeta = \pm\delta$ with $\delta = 0.084$.




Fig. 3. Scans along $Q = (1/3, 1/3, \zeta)$ with (a) $\zeta = 0$ and (b) $\zeta = 6$. Solid lines are the results of Gaussian fitting.

The temperature dependence of the scan profile (see Fig. 4) shows that possible changes of δ is negligible, similar to pure CsCuCl_3 . The possibility that the central peak and the side peaks come from different magnetic domains is excluded, because the temperature variations of their intensities are strongly correlated as shown in Fig. 5. We see that the magnetic unit cell is enlarged by $\sqrt{3} \times \sqrt{3}$ in the basal plane, which is characteristic of the triangular spin arrangement. The side peaks indicate that the helical spin structure is realized along the c -axis. The value of δ corresponds to the period of the helix. The period is evaluated as $6/\delta = 71.4 \text{ Cu-sites}$, which is almost the same as that in CsCuCl_3 , i.e., 70.6 Cu-sites .¹⁰⁾

It is remarkable that in the present system, the magnetic Bragg peak is observed at $Q = (1/3, 1/3, 0)$, which




Fig. 4. Scans along the c^* -axis around $Q = (1/3, 1/3, 0)$ at various temperatures.




Fig. 5. Magnetic Bragg intensities at $Q = (1/3, 1/3, 0)$ and $(1/3, 1/3, 0.084)$ as a function of temperature. Néel temperatures are indicated by arrows.

is absent in CsCuCl_3 . On the other hand, around $(1/3, 1/3, 6)$, the central peak is very weak as compared with the side peaks (see Fig. 3). Since the angle between the scattering vector and the c^* -axis is 90° for $Q = (1/3, 1/3, 0)$ and 15.7° for $Q = (1/3, 1/3, 6)$, the c -axis component of spins is much less visible around the $(1/3, 1/3, 6)$ reflection than around the $(1/3, 1/3, 0)$ reflection. On the other hand, the basal plane component

of spins make a larger contribution to the magnetic reflections around $(1/3, 1/3, 6)$. Thus, it is deduced that the central peaks originate from the c -axis component of spins, while the basal plane component gives rise to the side peaks as for CsCuCl_3 . We refer to a plane spanned by the spins on a triangular lattice as *spin-plane* in this paper. The present result leads us straightforward to conclude that the phase II is an oblique triangular phase, in which the spin-plane is tilted with an angle ϕ from the basal plane (see Fig. 6 (a) and (b)). The spin-planes are




Fig. 6. (a) Oblique triangular antiferromagnetic structure in the low-temperature phase. Tilted spin planes represented by circles are stacked along the c -axis with forming a helix with a pitch 5 degrees. (b) The angle ϕ indicates the tilt between the spin plane and the c -plane. (c) The angle γ is half of the angle between S_2 and S_3 .

stacked along the c -axis with forming a helix with a pitch $360 \times \delta/6 \sim 5$ degrees.

It is quite natural to consider that the spin structure in each spin-plane is close to the 120° -structure, because there is no anisotropy which is strong enough to deform largely the 120° -structure or lift the moments from the spin-plane. Although the D-M interaction is of the same order in magnitude as J_1 , its effect is reduced to be $\sim D^2/J_0$ due to the competition with the intrachain exchange interaction J_0 .⁹⁾ In order to take account of a slight deformation from the 120° -structure, the angle 2γ between the spins S_2 and S_3 , is adopted as a variable parameter as illustrated in Fig. 6 (c). Under the assumption that the thermal averages of spins on all sublattices are the same, the integrated intensity of the magnetic Bragg peaks for $Q = (h, h, \zeta)$ with $h = n/3$ is given by

$$I \propto |f(Q)|^2 L(\theta) |F(h, h, \zeta)|^2 \quad (3.1)$$

$$L(\theta) = \cos \left(\frac{\pi}{2} - \theta - \arccos \sqrt{\frac{a^2 \zeta^2}{4c^2 h^2 + a^2 \zeta^2}} \right)$$

and $I = 0$ for integer h , where θ is half of the scattering angle, $L(\theta)$ is the Lorentz factor for the parallel scans, $f(Q)$ is the magnetic form factor and a and c are lattice constants. The values of $f(Q)$ were taken from reference.¹⁹⁾ $F(h, h, \zeta)$ is the magnetic structure factor. Due to the long-period helical spin structure along the c -axis, the component in the a - b -plane and the c -axis component of the spins give Bragg peaks at $\zeta = \pm\delta$ and $\zeta = 0$, respectively. The structure factors for the planar component F_{ab} and the axial component F_c are described as

$$|F_{ab}(hh\zeta)|^2 = d_{(hh\zeta)}^2 \left\{ 2 \left(\frac{h}{a} \right)^2 + \left(\frac{\zeta}{c} \right)^2 \right\} \times \{ (1 + \cos \gamma)^2 \cos^2 \phi + 3 \sin^2 \gamma \} \quad (3.2)$$

$$|F_c(hh\zeta)|^2 = d_{(hh\zeta)}^2 \left(\frac{2h}{a} \right)^2 (1 + \cos^2 \gamma) \sin^2 \phi, \quad (3.3)$$

where $d_{(hh\zeta)}$ is the spacing of the (h, h, ζ) lattice plane. Integrated Bragg intensities are calculated for various angles of ϕ and γ . In Fig. 7 the variation of the reliability factor R given by

$$R = \sum_{h,h,\zeta} |I_{\text{calc}}(h, h, \zeta) - I_{\text{obs}}(h, h, \zeta)| / \sum_{h,h,\zeta} I_{\text{obs}}(h, h, \zeta).$$

is represented as a contour map. We see that the R -factor




Fig. 7. Contour plot of the reliability factor R as a function of the angle γ and ϕ . Dot dashed line indicates the minimized R -factor $R = 0.182$. Dashed line indicates the line which satisfies eq. (3.5).

has the same minimum value on the dot-dashed line in Fig. 7. With decreasing tilting angle ϕ , the canting angle γ which gives minimum R -factor rapidly decreases, and become zero when $\phi = 29.2^\circ$. Since the magnetic structure could not be determined uniquely only from the magnetic Bragg intensities, the value of the spontaneous magnetization along the c -axis detected from magnetization measurements (see Fig. 2 on ref.¹⁸⁾) is necessary to give the conclusive answer. For the oblique triangular

structure, each sublattice spin is described as

$$\begin{aligned} S_1 &= S(\cos \phi, 0, -\sin \phi) \\ S_2 &= S(-\cos \gamma \cos \phi, -\sin \gamma, \cos \gamma \sin \phi) \\ S_3 &= S(-\cos \gamma \cos \phi, \sin \gamma, \cos \gamma \sin \phi). \end{aligned} \quad (3.4)$$

Hence, the spontaneous magnetization M_{sp} per site is given by

$$M_{\text{sp}} = \frac{2 \cos \gamma - 1}{3} \sin \phi [\mu_B / \text{magnetic ion}]. \quad (3.5)$$

In Fig. 7, the dashed line gives $M_{\text{sp}} = 0.025 \mu_B$ which was observed by the previous magnetization measurements.¹⁸⁾ From the intersection of dashed and dot-dashed lines, the angles ϕ and γ are obtained as $(\phi, \gamma) = (42^\circ, 56^\circ)$.

In Table I, we show the experimental and calculated intensities of the Bragg peaks around $Q = (1/3, 1/3, 0)$, $(1/3, 1/3, 6)$ and $(2/3, 2/3, 0)$ together with the calculated intensities which corresponds to the A- and B-point in Fig. 7. Experimental and calculated intensities are normalized for the total intensities of central and side peaks around $(1/3, 1/3, 0)$. The spin structure of CsCuCl_3 correspond to the case $\phi = 0^\circ$ and $\gamma = 60^\circ$ which is indicated by A in Fig. 7. We see from Table I that the spins form the triangular structure neither in the c -plane, nor in the plane including the c -axis, but in a plane tilted from the c -plane with the angle $\phi = 42^\circ$ and that the triangular spin structure ($\gamma = 56^\circ$), is close to the 120° spin structure. The difference between the observed and calculated values of the scattering intensities may come from the fact that we did not taking into account of the variation of the magnitude of S for each sublattice.

As shown in Fig. 5, with increasing temperature, the intensity of the central peak decreases and disappears at $T_{\text{N}2} = 7.2$ K. On the other hand, the intensity of the side peak is almost constant up to $T_{\text{N}2}$, and decreases rapidly, and becomes zero at $T_{\text{N}1} = 9.5$ K. Both phase transitions look sharp, which indicates a good homogeneity of the sample. From the present result, we see that the tilting angle ϕ decreases with increasing temperature, and becomes zero at $T_{\text{N}2}$. Thus, we can deduce that the phase I between $T_{\text{N}2}$ and $T_{\text{N}1}$ is identical to the ordered phase of CsCuCl_3 , which is consistent with the fact that the Co^{2+} doping produces the phase II in the low-temperature and low-field region in the ordered state of CsCuCl_3 , and its area increases with increasing the Co^{2+} concentration x .

CsCuCl_3 has a weak planar anisotropy, while CsCoCl_3 is an Ising spin system.¹⁶⁾ Phase transitions in the random spin systems with competing anisotropies have been investigated in $\text{Fe}_{1-x}\text{Co}_x\text{Cl}_2$ ²⁰⁾ and $\text{Fe}_{1-x}\text{Co}_x\text{Cl}_2 \cdot 2\text{H}_2\text{O}$ ²¹⁾ by means of neutron scattering. Both systems undergo two phase transitions, which are characterized by the orderings of two spin components. In $\text{Fe}_{1-x}\text{Co}_x\text{Cl}_2$, the ordering of one spin component is drastically altered by the ordering of the other component, so that the lower-temperature transition is rather smeared. On the other hand, in $\text{Fe}_{1-x}\text{Co}_x\text{Cl}_2 \cdot 2\text{H}_2\text{O}$, two phase transitions are sharp, and two order parameters are decoupled. $\text{CsCu}_{0.97}\text{Co}_{0.03}\text{Cl}_3$ differs from the both in the ordering nature. In the

Table I. Experimental and calculated intensities of several magnetic Bragg reflections in $\text{CsCu}_{1-x}\text{Co}_x\text{Cl}_3$ with $x = 0.03$ at $T = 1.6$ K. They are normalized to the total peak intensities around $Q = (1/3, 1/3, 0)$ reflection. The calculated intensities at A- and B-point in Fig. 7 are also shown for comparison.

(h, h, ζ)	I_{obs}	$I_{\text{cal}} \left(\begin{smallmatrix} \phi = 42^\circ \\ \gamma = 57^\circ \end{smallmatrix} \right)$	$I_{\text{cal}}(\text{A}) \left(\begin{smallmatrix} \phi = 0^\circ \\ \gamma = 60^\circ \end{smallmatrix} \right)$	$I_{\text{cal}}(\text{B}) \left(\begin{smallmatrix} \phi = 90^\circ \\ \gamma = 57.49^\circ \end{smallmatrix} \right)$
$(\frac{1}{3}, \frac{1}{3}, -\delta)$	0.324	0.306	0.497	0.155
$(\frac{1}{3}, \frac{1}{3}, 0)$	0.384	0.384	0	0.689
$(\frac{1}{3}, \frac{1}{3}, \delta)$	0.292	0.310	0.503	0.156
$(\frac{1}{3}, \frac{1}{3}, 6 - \delta)$	0.210	0.230	0.373	0.116
$(\frac{1}{3}, \frac{1}{3}, 6)$	0.012	0.011	0	0.019
$(\frac{1}{3}, \frac{1}{3}, 6 + \delta)$	0.186	0.226	0.366	0.114
$(\frac{2}{3}, \frac{2}{3}, -\delta)$	0.171	0.266	0.432	0.135
$(\frac{2}{3}, \frac{2}{3}, 0)$	0.252	0.330	0	0.573
$(\frac{2}{3}, \frac{2}{3}, +\delta)$	0.172	0.266	0.432	0.134
		$R = 0.182$	$R = 0.947$	$R = 0.598$
$M_{\text{sp}} [\mu_{\text{B}}/\text{ion}]$	0.025	0.025	0	0.025

present system, the basal plane component of spins orders first at T_{N1} and then the c -axis component orders at T_{N2} , so that the oblique triangular spin structure is realized. Both phase transitions are fairly sharp, and both components are not decoupled, *i.e.*, the growth of the basal plane component of spins below T_{N2} is suppressed due to the growth of the c -axis component which is equivalent to the increase of tilting angle ϕ . The present system is the first example that undergoes the definite phase transition to the oblique spin phase. Its mechanism seems not to be explained by only the competing anisotropies, because mixed triangular antiferromagnetic systems with axial and planar anisotropies such as $\text{Rb}_{1-x}\text{K}_x\text{NiCl}_3$ ²²⁾ and $\text{CsMn}(\text{Br}_x\text{I}_{1-x})_3$ ^{23,24)} exhibited neither mixed ordered phase nor oblique phase.

In CsCuCl_3 , the off-diagonal exchange term of the form

$$\mathcal{H}_{\text{od}} = J_{ij}^{zx}(S_i^z S_j^x + S_i^x S_j^z) + J_{ij}^{zy}(S_i^z S_j^y + S_i^y S_j^z) \quad (3.6)$$

is allowed, because the elongated axes of CuCl_6 octahedra are canted from the c -axis, so that the local symmetry is lower than the trigonal one. Here we take the z -axis parallel to the c -axis and x - and y -axes in the basal plane. The off-diagonal exchange term has been proposed to interpret the nature of the mixed ordered state in $\text{Fe}_{1-x}\text{Co}_x\text{Cl}_2$.^{20,25)} Recently, Pleimling²⁶⁾ discussed the ordering of the perpendicular spin component in the Ising-like system FeBr_2 on the basis of the off-diagonal exchange term. Due to the off-diagonal term, the ordering of the basal plane component of spins can induce the ordering of the c -axis component. We suggest that the doping of small amounts of Co^{2+} ions suppresses the planar anisotropy in the average, so that the effect of the off-diagonal term is relatively enhanced. However, whether the orderings of both components occur simultaneously or separately as observed in the present system is unclear. This problem should be investigated theoret-

ically.

§4. Conclusion

We have performed elastic neutron scattering experiments on the triangular antiferromagnetic system $\text{CsCu}_{0.97}\text{Co}_{0.03}\text{Cl}_3$ which undergoes two phase transitions at $T_{\text{N1}} = 9.5$ K and $T_{\text{N2}} = 7.2$ K. It was found that the low-temperature phase is an oblique triangular antiferromagnetic phase with the spin-plane tilted from the basal plane, and that the intermediate phase is identical to the ordered phase of CsCuCl_3 . We suggest that the off-diagonal exchange interaction gives rise to the oblique phase in the present system.

Acknowledgments

The authors wish to thank K. D. Schotte and H. Shiba for useful discussions.

- 1) H. Nojiri, K. Takahashi, T. Fukuda, M. Fujita, M. Arai and M. Motokawa: *Physica B* **241-243** 210 (1998).
- 2) T. Werner, H. B. Weber, J. Wosnitza, A. Kelnberger, M. Meschke, U. Schotte, N. Stüßer, Y. Ding and M. Winkelmann: *Solid State Comm.* **102** 609 (1997).
- 3) U. Schotte, A. Kelnberger and N. Stüßer: *J. Phys.: Condens. Matter* **10** 6391 (1998).
- 4) A. E. Jacobs and T. Nikuni: *J. Phys.: Condens. Matter* **10** 6405 (1998).
- 5) A. W. Schlueter, R. A. Jacobson and R. E. Rundle, *Inorg. Chem.* **5** 277 (1966).
- 6) C. J. Kroese, W. J. A. Maaskant and G. D. Verschoor, *Acta Crystallogr.* **B30** 1053 (1974).
- 7) S. Hirotsu, *J. Phys.* **C10** 967 (1977).
- 8) Y. Tazuke, H. Tanaka, K. Iio and K. Nagata: *J. Phys. Soc. Jpn.* **50** 3919 (1981).
- 9) H. Tanaka, U. Schotte and K. D. Schotte, *J. Phys. Soc. Jpn.* **61** 1344 (1992).
- 10) K. Adachi, N. Achiwa and M. Mekata: *J. Phys. Soc. Jpn.* **49** 545 (1980).
- 11) H. B. Weber, T. Werner, J. Wosnitza, H. von Löhneysen and U. Schotte, *Phys. Rev. B* **54** 15 924 (1996).

- 12) T. Nikuni and H. Shiba, J. Phys. Soc. Jpn. **62** 3268 (1993).
- 13) H. Nishimori and S. Miyashita, J. Phys. Soc. Jpn. **64** 4448 (1986).
- 14) A. V. Chubukov and D. I. Golosov: J. Phys.: Condens. Matter **3** 69 (1991).
- 15) A. Honecker, M. Kaulke and K. D. Schotte: Euro. Phys. J. **B15** 423 (1999).
- 16) M. F. Collins and O. A. Petrenko: Can. J. Phys. **75** 605 (1997) and references therein.
- 17) M. Mekata and K. Adachi: J. Phys. Soc. Jpn. **44** 806 (1978).
- 18) T. Ono, H. Horai and H. Tanaka: J. Phys.: Condens. Matter **12** (2000) 975.
- 19) R. E. Watson and A. J. Freeman: Acta Crystallogr. **14** (1961) 27.
- 20) P. Wong, P. M. Horn, R. J. Birgeneau and G. Shirane: Phys. Rev. **B27** 428 (1983).
- 21) K. Katsumata, H. Yoshizawa, G. Shirane and R. J. Birgeneau: Phys. Rev. **B27** 316 (1985).
- 22) H. Tanaka, T. Hasegawa and K. Nagata: J. Phys. Soc. Jpn. **62** 4053 (1993).
- 23) T. Ono, H. Tanaka, T. Kato and K. Iio: J. Phys.: Condens. Matter **10** 7209 (1998).
- 24) T. Ono, H. Tanaka, T. Kato, K. Nakajima and K. Kakurai: J. Phys.: Condens. Matter **11** 4427 (1999).
- 25) D. Mukamel: Phys. Rev. Lett. **46** 845 (1981).
- 26) M. Pleimling: Eur. Phys. J. **B10** 465 (1999).